



EOSDIS

NASA'S EARTH OBSERVING SYSTEM
DATA AND INFORMATION SYSTEM

Using UMM-Var and E2E To Improve The User Experience for Accessing NASA EOSDIS Data Sets

ESIP Summer Meeting 2019

Tyler Stevens
EED-2 Senior Discipline Engineer
Tyler.B.Stevens@nasa.gov

Simon Cantrell
EED-2 Senior Principal Systems Engineer
simon.cantrell@nasa.gov

This work was supported by NASA/GSFC under Raytheon Co. contract number NNG15HZ39C.
This document does not contain technology or Technical Data controlled under either the U.S. International Traffic
in Arms Regulations or the U.S. Export Administration Regulations.

Outline

- I. Introduction To UMM-Var
- II. E2E Capability and End-User Experience
- III. Demo of E2E Capability in Earthdata Search

I. Introduction To UMM- Var

What is the UMM-Var?

The Unified Metadata Model for Variables (UMM-Var) describes variable-level metadata. UMM-Var provides a higher level of specificity about the data than collections, and allows search and discovery of data at the variable level using community-sourced terms.

- Enables services at the variable level, as an extension to the UMM-C (Collections) and UMM-G (Granules) metadata models.
- Permits users to search and discover variables via well known geophysical terms (a.k.a. measurement names), as well as using existing Science Keywords.

Measurement: An observable property, usually geophysical. For models, it is a simulated observable property. Measurement Names are of the form: <object>__<quantity>. Names shall contain only lowercase letters and numbers along with the Standard Names separator characters (__, -, ~, __).

Examples:

"land_surface_air_flow",
"land_surface_air_heat-incoming-latent",
"land_surface_air-incoming-sensible",


Variable: A measured geophysical quantity associated with a particular data product, typically a point, swath or grid of values nested within the data file. One or more variables may be associated with a given collection.

Examples:

psl (mean sea level pressure)
O3_ppbv (ozone mixing ratio reported in parts per billion by volume)
Scat_550 (total dry aerosol scattering coefficient at 550 nm)

Fields in UMM-Var

This view highlights the top-level fields in the UMM-Var.



What Does a Provider Need to Know About UMM-Var?

- **UMM-Var Version 1.5**

- Updated the use case for Size Estimation Service.
- Updated to support Size Estimation Service with better specification of data formats (via an enumeration) and a few fixes.
- Improvements following the Goddard Earth Sciences (GES) Data and Information Services Center (DISC) pathfinder work resulted in a refinement of the implementation of Variable uniqueness; added AcquisitionSourceName.

- **UMM-Var Metadata Curation**

- Records can be curated using the UMM-Var Generation (UVG) script.
- Records can be curated using the Metadata Management Tool (MMT).
- Records can be curated using the Common Metadata Repository (CMR) API.
- Records can be associated to a related collection record.

- **UMM-Var Documentation**

- Access the model and associated documentation at <https://wiki.earthdata.nasa.gov/display/CMR/CMR+Documents>.

II. End-To-End Service (E2E) Capability and End- User Experience

End-to-End Services (E2E) Capability

- Enables users to apply data transformations to data sets at the variable level.
- Requires both UMM-S and UMM-Var records to be present in CMR.
 - A UMM-S record is associated with all relevant UMM-C records.
 - The Service Options class in UMM-S provides a means to define the data transformations options available via a specific service.
- In Earthdata search, the service options show up for each of the variables within a collection.
- E2E goes beyond the traditional “order and download” paradigm and offers a more enhanced way to retrieve data.

What Does a Provider Need To Do to Add E2E Capabilities To Their Records?

- **Update UMM-S Records**
 - Curate record using the MMT or CMR API.
 - Add ServiceOptions to services, showing data transformations available, e.g. via OPeNDAP, THREDDS, WMS services (Note: E2E currently supports OPeNDAP only).
 - Add service to collection associations via MMT or the CMR API.
- **Update UMM-Var Records**
 - Use the UVG script to generate all the needed variable records.
 - By default, UVG adds variable–collection associations via the CMR API at the time of variable generation.
 - Manually curate records post-generation using the MMT or CMR API (note: there are ongoing improvements to the UVG script based on provider testing).


E2E Documentation

- Existing collection with **new services**:
 - <https://wiki.earthdata.nasa.gov/x/OhDZBw>
- **New collection** capable of services:
 - <https://wiki.earthdata.nasa.gov/x/NRDZBw>
- How To: Configure Customizable Icons:
 - <https://wiki.earthdata.nasa.gov/x/9ZsYC>
- UMM-S Curators Guide
 - <https://wiki.earthdata.nasa.gov/x/agFHC>

User Experience in Earthdata Search

Collection Search

Search for collection in Earthdata Search (<https://search.uat.earthdata.nasa.gov/>).



Customizable Badge

The Customizable badge is displayed to highlight that spatial, variable, reformatting, and temporal subsetting options are available on a given collection. These options are set in the UMM-Var metadata.



User Experience in Earthdata Search

Project Page

You can customize your data access and subsetting options.

Edit Options
MERRA-2 tavg1_2d_slv_Nx: 2d,1-Hourly, Time-Averaged, Single-Level,Assimilation,Single-Level Diagnostics 0.625 x 0.5 degree V5.12.4 (M2T1NXSLV) at GES DISC

Select Data Access Method

☐ Direct Download ?

☒ Customize (OPeNDAP) ?

Variable Selection
Use science keywords to subset your collection granules by measurements and variables.
No variables selected. All variables will be included in download.

[Edit Variables](#)

Output Format Selection
Choose from output format options like GeoTIFF, NETCDF, and other file types.

ASCII

✓ Collection 1 of 1 [Done](#)

Variable Selection

Select specific variables for a given collection.

Customizations
MERRA-2 tavg1_2d_slv_Nx: 2d,1-Hourly, Time-Averaged, Single-Level,Assimilation,Single-Level Diagnostics 0.625 x 0.5 degree V5.12.4 (M2T1NXSLV) at GES DISC

[Back to Edit Options](#)

Variable Selection

[ALL LEAFNODES](#) ☒ AIR TEMPERATURE

☐ Select All Variables

PS
☐ surface_pressure
[View Details](#)

SLP
☐ sea_level_pressure
[View Details](#)

lat
☐ latitude
[View Details](#)

lon

[Back](#) [Save](#)

III. Demo of E2E Capability in Earthdata Search

Demo of E2E Capability in Earthdata Search

User Scenario:

I am looking for soil moisture data from SMAP in the Washington State region and want to reformat and download the data so I can use the data in a downloadable tool that supports HDF formatted data.

Thank You

- If you have any questions, please reach out to me (Tyler.B.Stevens@nasa.gov) or Valerie Dixon (Valerie.dixon@nasa.gov)

This work was supported by NASA/GSFC under
Raytheon Co. contract number NNG15HZ39C.

Raytheon

in partnership with

